



# Specticle® G

**Active Ingredients:** indaziflam

**Formulation:** granule

**WSSA Group:** 29

**Signal Word:** no signal word required

## Best Uses

- Granular pre-emergent herbicide controls the most troublesome grassy and broadleaf weeds, as well as annual sedges
- Extended residual control of annual bluegrass (*Poa annua*), goosegrass, crabgrass and more than 90 listed weeds

## Key Strengths

- Broad-spectrum pre-emergent herbicide
- Long residual
- Low use rates
- Broadleaf and grassy weed control

## Attributes Of Specticle® G

- Controls weeds by inhibiting cellulose biosynthesis – its primary site of action is the newly developing root of a germinated weed seedling
- No stains, no odor formulation
- Newer mode-of-action – controls weeds known to develop resistance to other herbicides
- Works in turf, landscapes and hardscapes



*Poa annua*

*Eleusine indica*

*Digitaria sanguinalis*

## How To Apply

- Apply before weed seeds germinate
- Use as a broadcast treatment on bare ground or mulched areas around labeled ornamental plants and grasses
- Specticle G performs best when irrigation or rainfall occurs within 48 hours after application
- Adequate soil moisture is needed to activate Specticle G and provide maximum control



# Spect(i)cle® G

## Use Rates For Single Application Of Specticle G

| Target Weed  | Specticle G Use Rate (lbs per acre)** | Specticle G Use Rate (lbs per 1,000 sq ft) |
|--|---------------------------------------|--|
| Crabgrass<br>Goosegrass<br>Annual bluegrass<br>Broadleaf weeds | 100 - 200                             | 2.3 - 4.6                                  |
| Annual sedge<br>Kyllinga                                       | 200                                   | 4.6  |

\*Use higher use rates within the rate range for each target weed when a site has historically higher weed pressure.

\*\*The maximum single application on turf is 200 lbs. per acre (4.6 lbs. per 1,000 sq. ft.). Do not exceed 400 lbs. of Specticle G per acre (9.2 lbs. per 1,000 sq. ft.) in a 12-month period.

## Use Rates For Split Application Of Specticle G

| Target Weed  | Specticle G Use Rate (initial application) |                       | Specticle G Use Rate (within 90 days of initial application) |                       |
|--|--|-----------------------|--|-----------------------|
|  | (lbs per acre)**                           | (lbs per 1,000 sq ft) | (lbs per acre)**   | (lbs per 1,000 sq ft) |
| Crabgrass<br>Goosegrass<br>Annual bluegrass<br>Broadleaf weeds | 100 - 200                                  | 2.3 - 4.6             | 100 - 200  | 2.3 - 4.6             |
| Annual sedge<br>Kyllinga                                       | 200  | 4.6                   | 125 - 200  | 2.9 - 4.6             |



goosegrass



large crabgrass



spotted spurge



Florida pusley



yellow woodsorrel

BackedByBayer.com | @BayerGolf | | Available on iTunes and Google play: BackedByBayer

ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS CAREFULLY.  
Bayer CropScience LP, Environmental Science Division, 2 TW Alexander Drive, Research Triangle Park, NC 27709. 1-800-331-2867. www.BackedByBayer.com.  
Bayer, the Bayer Cross and Specticle are registered trademarks of Bayer. Not all products are registered in all states. ©2015 Bayer CropScience LP. 14BPG-052-P11

Backed by  
**BAYER**